

real-world practice: the AMC Single Centre Real World PCI Registry. *EuroIntervention* 2015;**10**:1160–1168.

13. Verheye S, Ormiston JA, Stewart J, Webster M, Sanidas E, Costa R, Costa JR Jr, Chamie D, Abizaid AS, Pinto I, Morrison L, Toyloy S, Bhat V, Yan J, Abizaid A. A next-generation bioresorbable coronary scaffold system: from bench to first clinical evaluation: 6- and 12-month clinical and multimodality imaging results. *JACC Cardiovasc Interv* 2014;**7**:89–99.
14. Maeng M, Jensen LO, Falk E, Andersen HR, Thuesen L. Negative vascular remodelling after implantation of bioabsorbable magnesium alloy stents in porcine coronary arteries: a randomised comparison with bare-metal and sirolimus-eluting stents. *Heart* 2009;**95**:241–246.

15. Erbel R, Di MC, Bartunek J, Bonnier J, de Bruyne B, Eberli FR, Erne P, Haude M, Heublein B, Horrigan M, Ilesley C, Bose D, Koolen J, Luscher TF, Weissman N, Waksman R. Temporary scaffolding of coronary arteries with bioabsorbable magnesium stents: a prospective, non-randomised multicentre trial. *Lancet* 2007;**369**:1869–1875.
16. Haude M, Erbel R, Erne P, Verheye S, Degen H, Bose D, Vermeersch P, Wijnbergen I, Weissman N, Prati F, Waksman R, Koolen J. Safety and performance of the drug-eluting absorbable metal scaffold (DREAMS) in patients with de-novo coronary lesions: 12 month results of the prospective, multicentre, first-in-man BIOSOLVE-I trial. *Lancet* 2013;**381**:836–844.

## CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehv364

Online publish-ahead-of-print 10 August 2015

# Ultra-low-dose hybrid single photon emission computed tomography and coronary computed tomography angiography: a comprehensive and non-invasive diagnostic workup of suspected coronary artery disease

Dominik C. Benz<sup>1</sup>, Christian Templin<sup>2</sup>, Philipp A. Kaufmann<sup>1</sup>, and Ronny R. Buechel<sup>1\*</sup>

<sup>1</sup>Department of Nuclear Medicine, Cardiac Imaging, University Hospital Zurich, Ramistrasse 100, Zurich CH-8091, Switzerland; and <sup>2</sup>Department of Cardiology, University Hospital Zurich, Zurich, Switzerland

\* Corresponding author. Tel: +41 44 255 10 59, Fax: +41 44 255 44 28, Email: [ronny.buechel@usz.ch](mailto:ronny.buechel@usz.ch)

This article was guest edited by Brahmajee K. Nallamothu, University of Michigan, [bnallamo@umich.edu](mailto:bnallamo@umich.edu)

A 52-year-old male patient (body mass index: 23.5 kg/m<sup>2</sup>) without known coronary artery disease (CAD) or any cardiovascular risk factors was referred for non-invasive evaluation due to typical angina. The initial computed tomography (CT) for coronary artery calcium scoring (CACS) showed no calcifications. Subsequently, coronary computed tomography angiography (CCTA) was performed using prospective electrocardiogram triggering on a latest generation 256-slice CT scanner (Revolution CT, GE Healthcare). Coronary computed tomography angiography revealed a non-calcified lesion with subtotal occlusion in the middle left anterior descending (LAD) coronary artery (Panel A). Adenosine stress and rest single photon emission computed tomography (SPECT) myocardial perfusion imaging (MPI)—with administration of 143 and 276 Mbq <sup>99m</sup>Techetium-tetrofosmine for stress and rest imaging, respectively—was performed on a high-sensitive cadmium-zinc-telluride detector gamma camera (Discovery 530NM, GE Healthcare) for assessment of ischaemia or scar (Panel B stress; Panel C rest). Hybrid SPECT/CCTA demonstrated a large reversible perfusion defect in the apical and anteroseptal myocardium, subtended by the LAD (Panel D). Invasive coronary angiography confirmed the morphological finding (Panel E), and the patient was successfully revascularized.

Cumulative effective radiation dose for cardiac hybrid imaging (CACS 0.66 mSv, CCTA 0.47 mSv, SPECT 2.8 mSv) resulted in 3.93 mSv and total amount of contrast agent used for CCTA was 30 mL.

While adding SPECT MPI to CCTA has been demonstrated to improve diagnostic accuracy, the possible risks of radiation exposure from hybrid SPECT/CCTA imaging have remained an issue of concern. This case illustrates the feasibility of a comprehensive morphological and functional diagnostic work-up by cardiac hybrid SPECT/CCTA imaging of a patient presenting with stable CAD at a cumulative radiation dose <4 mSv.

Supplementary material is available at *European Heart Journal* online.

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author 2015. For permissions please email: [journals.permissions@oup.com](mailto:journals.permissions@oup.com).

